

II EU Workshop on Water Cherenkov Experiments  
for Precision Physics, Sept 2025


# Neutrinoless double beta decay search in KamLAND-Zen

Patrick Decowski  
[decowski@nikhef.nl](mailto:decowski@nikhef.nl)



UNIVERSITEIT VAN AMSTERDAM


# Double Beta Decay



$^{136}\text{Xe}$

A second-order process only detectable if first-order beta decay is energetically forbidden

Rare, but Standard Model Process:



$^{76}\text{Ge}$


$^{82}\text{Se}$

$^{100}\text{Mo}$

$^{130}\text{Te}$

...

# Neutrinoless Double Beta Decay



Is  $\nu$  Majorana?

$$M_\nu \neq 0$$


$$|\Delta L| = 2$$

$$0\nu\beta\beta : (A, Z) \rightarrow (A, Z + 2) + e^- + e^-$$

- Extremely rare radioactive process
- Requires massive Majorana neutrino
- Lepton Number Violation
- Model dependent - Standard interpretation: light Majorana  $\nu$  + SM interactions
  - Measure of neutrino mass scale  $\rightarrow$  effective Majorana mass  $\langle m_{\beta\beta} \rangle$

# Detecting $0\nu 2\beta$ Decay

Without energy resolution



Searching for this peak  
to measure  $T_{1/2}$


$0\nu\beta\beta$

$$\sum E_e/Q$$

In  $^{136}\text{Xe}$ :  
 $Q_{\beta\beta} = 2.458 \text{ MeV}$

# Detecting $0\nu 2\beta$ Decay

**With** energy resolution



Searching for this peak  
to measure  $T_{1/2}$

In  $^{136}\text{Xe}$ :  
 $Q_{\beta\beta} = 2.458 \text{ MeV}$

# What mass does $0\nu\beta\beta$ measure?

$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu}(Q, Z) |M_{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$

# What mass does $0\nu\beta\beta$ measure?

$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu}(Q, Z) |M_{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$

Phase Space factor:  
Calculable

Nuclear Matrix Element:  
Hard to calculate

# What mass does $0\nu\beta\beta$ measure?

$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu}(Q, Z) |M_{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$

Phase Space factor:  
Calculable


Nuclear Matrix Element:  
Hard to calculate

Interesting physics

Effective Majorana mass:  $\langle m_{\beta\beta} \rangle = \left| \sum_{i=1}^3 U_{ei}^2 m_i \right|$  [coherent sum]


"Inverted Ordering" :  $m_{\nu_1} > m_{\nu_3}$

"Normal Ordering" :  $m_{\nu_3} > m_{\nu_1}$





# Neutrino Science with KamLAND

Highly versatile KamLAND detector allows for a broad science program...






# KamLAND-Zen Collaboration



# KamLAND(-Zen) detector


Located in the  
Kamioka Mine in Japan  
1000m rock = 2700 mwe

- 1 kton Liquid Scintillator Detector
  - 6.5m radius balloon filled with:
    - 20% Pseudocumene (scintillator)
    - 80% Dodecane (oil)
    - PPO
- 34% PMT coverage
  - ~1300 17" fast PMTs
  - ~550 20" large PMTs
- Water Cherenkov veto
- Operational since 2002



# KamLAND(-Zen) detector

Particles interact in the LS and deposit energy.  
Energy is converted to light and detected by PMTs




Energy reconstruction

$$\sim \frac{6.7\%}{\sqrt{E(\text{MeV})}}$$

Position reconstruction

$$\sim \frac{13.7 \text{ cm}}{\sqrt{E(\text{MeV})}}$$




# KamLAND-Zen 400

2011 - 2015

- Inner-balloon radius 1.54m
- 320 - 380kg  $^{136}\text{Xe}$
- Exposure 504 kg-yr

$$T^{0\nu}_{1/2} > 1.1 \times 10^{26} \text{ yr}$$




KamLAND-Zen Coll, Phys. Rev. Lett. 117, 082503 (2016); arXiv:1605.02889

# KamLAND-Zen 800

2019 - 2024

- Inner-balloon radius 1.90m
- 745kg  $^{136}\text{Xe}$
- Exposure 2100 kg-yr

Full dataset - this talk





KamLAND-Zen Coll, arXiv:2406.11438

# Signals and Backgrounds


- Expected signal: **peak for  $^{136}\text{Xe}$  at  $Q_{\beta\beta} = 2.458 \text{ MeV}$**
- Define Region of Interest (ROI) between 2.35-2.70 MeV
- Primary Backgrounds:

- $2\nu\beta\beta$  decays
- Cosmic muon spallation
- Radioactive contamination, e.g.  $^{214}\text{Bi}$
- Solar neutrinos



# KLZ 800 Inner-Ballon Backgrounds

$^{214}\text{Bi}$  from radon-chain veto:  $\beta$  delayed coincidence  $\alpha$  very effective veto in Xe-LS  
 But not on the balloon -  $\alpha$  may get absorbed by the balloon



Balloon film backgrounds:

$^{238}\text{U} \sim 4 \times 10^{-12}$  g/g


$^{232}\text{Th} \sim 2 \times 10^{-11}$  g/g

Xe-LS backgrounds:

$^{238}\text{U} \sim 1.5 \pm 0.4 \times 10^{-17}$  g/g

$^{232}\text{Th} \sim 3.0 \pm 0.4 \times 10^{-16}$  g/g

**10x reduction compared to KLZ 400 IB**




GEANT4 based MC with  $^{214}\text{Bi}$   $\beta+\gamma$  cascade, particle tracking, energy deposit, scintillation photon emission / propagation

# Muon Spallation

Carbon-based liquid scintillator:  $^{12}\text{C} + \mu \rightarrow$  spallation products


## Spallation on $^{12}\text{C}$



(muon time – decay time)

Triple coincidence cut (muon, neutron capture, subsequent  $^{10}\text{C}$  decay)  
effective veto, tagging efficiency  $\sim 99\%$

**Short-lived spallation products  $\tau < \sim 5$  min**





Isotope	$\tau$ (s)	Eff (%)
$^{10}\text{C}$	27.8	99.3
$^6\text{He}$	1.2	$97.6 \pm 1.7$
$^{137}\text{Xe}$	330	$74 \pm 7$

# Muon Spallation

$\mu + {}^{136}\text{Xe}$  spallation products from FLUKA simulation

Spallation on  ${}^{136}\text{Xe}$





Isotope	$T_{1/2}$ (s)
${}^{88}\text{Y}$	$9.2 \times 10^6$
${}^{124}\text{I}$	$3.6 \times 10^5$
${}^{130}\text{I}$	$4.5 \times 10^4$
${}^{110}\text{In}$	$1.8 \times 10^4$
${}^{132}\text{I}$	$8.3 \times 10^3$
${}^{118}\text{Sb}$	$2.2 \times 10^2$
${}^{122}\text{I}$	$2.2 \times 10^2$

# Muon Spallation

$\mu + {}^{136}\text{Xe}$  spallation products from FLUKA simulation, spectrum after decay

Spallation on  ${}^{136}\text{Xe}$



Isotope	$T_{1/2}$ (s)
${}^{88}\text{Y}$	$9.2 \times 10^6$
${}^{124}\text{I}$	$3.6 \times 10^5$
${}^{130}\text{I}$	$4.5 \times 10^4$
${}^{110}\text{In}$	$1.8 \times 10^4$
${}^{132}\text{I}$	$8.3 \times 10^3$
${}^{118}\text{Sb}$	$2.2 \times 10^2$
${}^{122}\text{I}$	$2.2 \times 10^2$

Long-lived spallation products in the ROI

$T_{1/2}$ : **several hours to weeks**


Very low rate!

# Event Selection

- Event selection cuts:

- Events  $< 2.5\text{m}$  from center and  $> 0.7\text{m}$  away from bottom
- Events  $> 150\text{ms}$  after muons
- Radioactive decays by coincidence cut rejected
- $\bar{\nu}_e$  identified by coincidence cut rejected
- Poorly reconstructed events rejected
- Spallation cuts applied:
  - Short-lived spallation (e.g.  $^{10}\text{C}$ ) rejected
  - Long-lived (LL) spallation: tagged and untagged sample


$(745 \pm 3)\text{kg Xe}$ , 1131 day lifetime  $\rightarrow$  2100 kg-yr exposure



# Fitting the Data in Equal Volume Bins

Vertex distribution in the ROI overlaid on  $^{214}\text{Bi}$  MC

Beta-decay of  $^{214}\text{Bi}$  can also include a  $\gamma$  at 2.448 MeV




- Simultaneously fit 40 equal volume bins inside of  $R < 2.5$  m
- Inner region  $\rightarrow$  more sensitive to  $0\nu 2\beta$  decay
- Outer region  $\rightarrow$  more sensitive to backgrounds on inner-balloon film
- All parameters fitted simultaneously

# Fitting the Data in Equal Volume Bins

Vertex distribution in the ROI overlaid on  $^{214}\text{Bi}$  MC

Beta-decay of  $^{214}\text{Bi}$  can also include a  $\gamma$  at 2.448 MeV




- Simultaneously fit 40 equal volume bins inside of  $R < 2.5$  m
- Inner region  $\rightarrow$  more sensitive to  $0\nu 2\beta$  decay
- Outer region  $\rightarrow$  more sensitive to backgrounds on inner-balloon film
- All parameters fitted simultaneously

# Fitting the Data in Equal Volume Bins

Vertex distribution in the ROI overlaid on  $^{214}\text{Bi}$  MC

Beta-decay of  $^{214}\text{Bi}$  can also include a  $\gamma$  at 2.448 MeV




- Simultaneously fit 40 equal volume bins inside of  $R < 2.5$  m
- Inner region  $\rightarrow$  more sensitive to  $0\nu 2\beta$  decay
- Outer region  $\rightarrow$  more sensitive to backgrounds on inner-balloon film
- All parameters fitted simultaneously

# Fitting the Data in Equal Volume Bins

Vertex distribution in the ROI overlaid on  $^{214}\text{Bi}$  MC


Beta-decay of  $^{214}\text{Bi}$  can also include a  $\gamma$  at 2.448 MeV



- Simultaneously fit 40 equal volume bins inside of  $R < 2.5$  m
- Inner region  $\rightarrow$  more sensitive to  $0\nu 2\beta$  decay
- Outer region  $\rightarrow$  more sensitive to backgrounds on inner-balloon film
- All parameters fitted simultaneously


# New KamLAND-Zen 800 Results

Spectrum inside R<1.57m Fiducial Volume




# New KamLAND-Zen 800 Results

Spectrum inside R<1.57m Fiducial Volume




# New KamLAND-Zen 800 Results

Spectrum inside R<1.57m Fiducial Volume



# New KamLAND-Zen 800 Results

Spectrum inside R<1.57m Fiducial Volume




90% C.L. Upper Limit < 10.0 events →  $T_{1/2} > 3.4 \times 10^{26}$  yr  
[Sensitivity  $T_{1/2} > 2.3 \times 10^{26}$  yr]



Combined with earlier KamLAND-Zen 400 results:  $T_{1/2} > 3.8 \times 10^{26}$  years

# $0\nu 2\beta$ and tagged Long-lived spectrum

$0\nu 2\beta$  candidates



Xe-spallation candidates




The two spectra ( $0\nu 2\beta$  and Xe-spallation) are fitted simultaneously to constrain the Xe-spallation BG

# Effective Majorana Mass

$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu}(Q, Z) |M_{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$

$$T_{1/2} > 3.8 \times 10^{26} \text{ yr}$$




# Effective Majorana Mass

$$(T_{1/2}^{0\nu})^{-1} = G_{0\nu}(Q, Z) |M_{0\nu}|^2 \langle m_{\beta\beta} \rangle^2$$


$$T_{1/2} > 3.8 \times 10^{26} \text{ yr}$$

$\langle m_{\beta\beta} \rangle$  excl. limit depends on  
Nuclear Matrix Elements




**Most stringent limit on**  $\langle m_{\beta\beta} \rangle$

For some NMEs we probe  
Inverted Ordering  $\nu$ -mass





# Transition to KamLAND2-Zen in progress!



KamLAND(-Zen)  
2002-2024


**R.I.P.**



KamLAND2-Zen  
Start in 2028

# KamLAND2-Zen

Design sensitivity of  $T_{1/2} > 2 \times 10^{27}$  yrs and  $\langle m_{\beta\beta} \rangle \sim 20$  meV





- Improved energy resolution: Winston Cones (x1.8), High-QE PMTs (x1.9)
  - 4% → 2% (x100 reduction in  $2\nu\beta\beta$  BG rate)
- Better tag long-lived spallation BG: State-of-the-art electronics
- Reduce BG on inner balloon: scintillating balloon

# KamLAND2-Zen

Design sensitivity of  $T_{1/2} > 2 \times 10^{27}$  yrs and  $\langle m_{\beta\beta} \rangle \sim 20$  meV



- Improved energy resolution: Winston Cones (x1.8), High-QE PMTs (x1.9)
  - 4%  $\rightarrow$  2% (x100 reduction in  $2\nu\beta\beta$  BG rate)
- Better tag long-lived spallation BG: State-of-the-art electronics
- Reduce BG on inner balloon: scintillating balloon



# KamLAND2-Zen

Design sensitivity of  $T_{1/2} > 2 \times 10^{27}$  yrs and  $\langle m_{\beta\beta} \rangle \sim 20$  meV



- Improved energy resolution: Winston Cones (x1.8), High-QE PMTs (x1.9)
  - 4% → 2% (x100 reduction in  $2\nu\beta\beta$  BG rate)
- Better tag long-lived spallation BG: State-of-the-art electronics
- Reduce BG on inner balloon: scintillating balloon



# KamLAND-Zen disassembly Fall'24 onwards




Calibration system, LS and Xe support systems disassembled




Empty detector, first look inside in ~25 yrs!




Outer Detector PMT disassembly



Empty Outer Detector



Empty Outer Detector



# Summary



- Neutrinoless double beta decay searches are the only practical method to search for Majorana neutrinos in a model-independent way
- All KamLAND-Zen data, 2.1 ton-year of exposure
- $T_{1/2}^{0\nu} > 3.8 \times 10^{26}$  yr (90% C.L.)  $\rightarrow \langle m_{\beta\beta} \rangle < 28 - 122$  meV
  - Currently best limit - starting to probe Inverted Ordering
  - KamLAND-Zen 800 stopped operation, being dismantled  $\rightarrow$  KamLAND2-Zen
- KamLAND2-Zen will have sensitivity of  $2 \times 10^{27}$  years  $\rightarrow \langle m_{\beta\beta} \rangle \sim 20$  meV


# KamLAND-Zen uses Xe-doped LS



- +Well-understood detector
- +Highly pure, self-shielding environment
- +Large  $\beta\beta$  source mass, scalable
- -Relatively poor energy resolution
- -No particle identification

$$T_{1/2}^{0\nu} \propto \epsilon \frac{a}{A} \sqrt{\frac{Mt}{b\Delta E}}$$

**Detector Mass,  
Exposure, BG and  
Energy Resolution**




745 kg of  $^{136}\text{Xe}$   
dissolved in Liquid  
Scintillator

# Sensitivity


KamLAND-Zen, 2.1 ton-yr exposure

Given the background, what is the expected sensitivity?



**Sensitivity:  $T_{1/2} > 2.6 \times 10^{26}$  years**  
**Combined limit:  $T_{1/2} > 3.8 \times 10^{26}$  years**


# KLZ 800 Inner-Ballon Backgrounds



Balloon film backgrounds:

$^{238}\text{U} \sim 3 \times 10^{-12} \text{ g/g}$   
 $^{232}\text{Th} \sim 4 \times 10^{-11} \text{ g/g}$

**10x reduction compared to KLZ 400 IB**



Rate dominated by  $2\nu 2\beta$

Xe-LS backgrounds:  
 $^{238}\text{U} \sim 1.5 \pm 0.4 \times 10^{-17} \text{ g/g}$   
 $^{232}\text{Th} \sim 3.0 \pm 0.4 \times 10^{-16} \text{ g/g}$

